Congratulations!

# 1<sup>st</sup> night @home Instructions

We take care of you!

On the next page you will find the most important intstructions for getting you through the 1<sup>st</sup> night.

(Click on the arrow)











Scan **ór** click the QR-code for website Babybalance!











Always call your midwife, if you have any questions wich can't wait till next morning!

<u>Mother</u>

**Baby** 

<u>Nutrition</u>

Bed/Hotwater bottles

<u>Temperature</u>

Changing baby



Click on the relevant topic in the bar above for more information and advice.







Always call your midwife, if you have any questions wich can't wait till next morning!

<u>Mother</u>

Baby


<u>Nutrition</u>

Bed/Hotwater bottles

<u>Temperature</u>

Changing baby

Wash your hands before every feeding moment and after every diaper change.









Always call your midwife, if you have any questions wich can't wait till next morning!

<u>Mother</u>

Baby

**Nutrition** 

Bed/Hotwater bottles

Temperature

Changing baby





- Initiate breastfeeding every 2 à 3 hours.
- Apply alternately (left/right) to the breast. You will start with the breast you finished with for the next feeding.
- Breastfeeding must not hurt! If so, then the baby doesn't have the nipple properly in their mouth.

#### Formula

- 1 leveled scoop of formula (powder) of 30 cc lukewarm water (±37°C).
- Measure water into the bottle.
- Heat water au-bain-marie or mix boiled water with cold water.
- Be aware that the nutrition in the bottle isn't too hot. Check this by dripping a drop of nutrition on the inside of your wrist.
- The baby may drink 10 to 20 cc every 3 hours.

Always call your midwife, if you have any questions wich can't wait till next morning!

<u>Mother</u>

Baby

**Nutrition** 

Bed/Hotwater bottles

Temperature

Changing baby







Never put a (waterproof) protection cover in the bed of the baby.



- Use a sheet and a blanket for making up the bed.
- Cover the bed briefly: baby's feet just barely touch the back of the bed, hydrophilic diaper under baby's head, double-folded blanket that reaches the chin.
- Baby lies on back in bed, turn baby's head after every feeding.

















Always call your midwife, if you have any questions wich can't wait till next morning!

<u>Mother</u>


Baby

<u>Nutrition</u>

Bed/Hotwater bottles

<u>Temperature</u>

Changing baby



Possibly: Warming up the baby skin to skin. Undress the baby, leave diaper and baby's hat on. Place the baby bare on the bare upper body of one of the parents, place a duvet or blanket over the baby.

- The correct temperature of the baby is between 36.5°C and 37.5°C.
- Make sure to temperature the baby at every feeding.
  - Temperature between 36.5 °C and 37.2 °C → put on a baby's hat on the baby, make a hot-water bottle and put safely in the bed with te baby.
  - o Temperature above 37.2 °C → keep baby's hat on, remove hot-water bottle from the bed.
  - o Temperature under 36.5 °C and above 37.5 °C → <u>call</u> <u>your midwife!</u>







Always call your midwife, if you have any questions wich can't wait till next morning!

<u>Mother</u>

**Baby** 

**Nutrition** 

Bed/Hotwater bottles

<u>Temperature</u>

Changing baby



### Special details by changing diaper:

- Black, 1<sup>st</sup>, baby's poop is difficult to remove.
   Suggestion! After cleaning baby's bum, apply ointment/Vaseline on baby's bum.
- Baby's urine can be orange/red, these are urates. If you're in doubt, you can save the diaper.
- Girls can have some slime from their vagina, possibly with some blood. It's called pre-menstrual, it's part of it en generally known.





**∢...** 

- Change baby's diaper by every feeding.
- Please record baby's temperature, feeding times, diapers (urine and poop) in the
- Special details by changing diaper.





Always call your midwife, if you have any questions wich can't wait till next morning!

Mother

Baby

**Nutrition** 

Bed/Hotwater bottles

<u>Temperature</u>

**C**'



- Don't go alone to the toilet/bathroom the 1st evening/night.

Tiredness and blood loss can make you dizzy.



- Urinate <u>before</u> every feeding + rinse with water.



- You must have urinated within **4 to 6** hours after birth of the placenta.



- You can lose blood clots, even up to fist size. Are you losing more than 2 large blood clots? Call the midwife!



Lot of blood loss, maternity pad is soaked within a half an hour ->
 call the midwife!



Always call your midwife, if you have any questions wich can't wait till next morning!

Mother

Baby

**Nutrition** 

Bed/Hotwater bottles

<u>Temperature</u>

Changing baby

Is hier het Engelstalig zorgplan gewenst en bijbehorend schema?

#### Instruction:

In your care plan you'll find this schedule. You can clearly record all the information for the maternity nurse.

Zorg

## Suggestion!

You can always save the used diapers, if you're in doubt.

Tijdstip	Geëindig met	Aantal cc gekolfd	Aantal cc bijvoeding	Aantal cc kunstvoeding	Urine	Ontlasting	Temp.	Opmerkingen
16	□ Links □ Rechts		.0		□ Ja □ Nee	□ Ja □ Nee		
	□ Links □ Rechts				□ Ja □ Nee	□ Ja □ Nee		
	□ Links □ Rechts		10		□ Ja □ Nee	□ Ja □ Nee		
	□ Links □ Rechts				□ Ja □ Nee	□ Ja □ Nee		
156	□ Links □ Rechts		.0		□ Ja □ Nee	□ Ja □ Nee		
	□ Links □ Rechts				□ Ja □ Nee	□ Ja □ Nee		